



CENTER FOR
ASTROPHYSICS
HARVARD & SMITHSONIAN

VERSION: 2.0
RELEASE DATE: September 17, 2025



**TROPOSPHERIC EMISSIONS:
MONITORING OF POLLUTION (TEMPO)
PROJECT**

**Total Ozone Level 2 and 3 Data products:
User Guide**

September 17, 2025

DOI: 10.5067/doc/tempo/O3TOT_L2-3UserGuide_V2.0

AUTHORS

Junsung Park (Center for Astrophysics | Harvard & Smithsonian)

Xiong Liu (Center for Astrophysics | Harvard & Smithsonian)

John Houck (Center for Astrophysics | Harvard & Smithsonian)

Dave Haffner (NASA Goddard Space Flight Center, Science Systems and Applications Inc.)

Kelly Chance (Center for Astrophysics | Harvard & Smithsonian)

CONTRIBUTIONS:

TEMPO Science Team, Center for Astrophysics | Harvard & Smithsonian

TEMPO Ground System Team, Center for Astrophysics | Harvard & Smithsonian

NASA Langley Research Center

NASA Goddard Space Flight Center

REVISION HISTORY

| Version | Item | Description | Release date |
|---------|--|---|--------------------|
| 1.0 | All | Initial version | May 20, 2024 |
| 1.1 | - Known data product issues - Sections 2, 3.2.2 and 4.3 | - Update known data product issues - Update validation status to provisional - Add a Section for data versioning and release history - Update quality flag explanation | December 9, 2024 |
| 2.0 | - Known data product issues - Sections 2 and 3 | - Update known data product issues - Update Algorithm overview - Update O3TOT usage recommendations | September 17, 2025 |

PRODUCT to SCIENCE DATA PROCESSING CENTER PIPELINE VERSION

| Version Designation | Science Data Processing Center Pipeline Version | Release |
|---------------------|---|--|
| V03 | 4.4 | May 20, 2024; first public release of TEMPO Level 2 and Level 3 products |
| V04 | 4.7 | September 17, 2025 |

KNOWN DATA PRODUCT ISSUES

Total Ozone (O3TOT)

- (1) TEMPO V03 O3TOT Total column ozone (TCO) shows latitude-dependent biases compared to OMI and OMPS TCO, varying from ~5% at lower latitudes to ~-2% at high latitudes. This latitudinal dependence does not vary much with season. This latitudinal dependence is confirmed from validation using ground-based Pandora, Dobson, and Brewer observations (Zhao et al., 2025). Preliminary testing indicates that this latitudinal dependence results mainly from the spectral dependence of radiometric bias in the Sun-normalized radiances. In addition, V03 O3TOT UV aerosol index shows negative biases of ~-2, primarily from the large overestimation in the normalized radiance of V03 L1b data.
 - a. This latitude dependence is caused by an issue in the TEMPO L1B data, which exhibits an additional bump feature in the irradiance spectrum between 300 nm and 330 nm after a wavelength-independent reduction of normalized radiance by ~13.4% in the UV band of V04 L1b data. To correct this, we implemented soft calibration at the wavelengths used in the ozone retrieval (312.61 nm and 317.62 nm). Additionally, we performed soft calibration to the wavelength (360.15 nm) used in the reflectance and aerosol index calculation.
 - b. After applying these soft calibrations after the L1b update, the latitude dependence has been mitigated in the O3TOT V04 data product. Preliminary evaluation indicates that the TCO from the TEMPO O3TOT algorithm shows excellent agreement, with a relative difference within $\pm 1\%$, with TCO from NPP OMPS satellite observations.
 - c. The peak of the histogram of UV aerosol index is improved from ~-2.5 to ~0, mainly from the L1b update and also slightly from additional soft calibrations.
- (2) The quality_flag has an issue (including unnecessary flags for the TEMPO observation) in the V03 TEMPO O3TOT product.
 - a. This issue has been updated from the V04 TEMPO O3TOT product. Therefore, users can confidently use good samples with quality_flag set to 0 (with $SZA \leq 80$, $VZA \leq 80$, and $CF \leq 0.5$).
- (3) TEMPO TCO also shows clear discontinuity when the algorithm switches from using the main pair of wavelengths (~317 and 331 nm) to a different wavelength pair (331 and 360 nm) at high solar zenith angle or viewing zenith angle, resulting in lower TCO values.
 - a. To address this, the thresholds for wavelength switch and profile shape change will be updated to reflect TEMPO characteristics. These thresholds will determine the appropriate values using a sufficiently large set of TEMPO O3TOT V04 product and incorporate them in the next product version.

Contents

| | |
|---|-----------|
| AUTHORS | 2 |
| REVISION HISTORY | 3 |
| PRODUCT to SCIENCE DATA PROCESSING CENTER PIPELINE VERSION | 3 |
| KNOWN DATA PRODUCT ISSUES | 4 |
| Total Ozone (O3TOT) | 4 |
| 1. Before You Begin..... | 6 |
| 1.1. Purpose of the Document..... | 6 |
| 1.2. Acronyms, Abbreviations, and Definitions | 6 |
| 2. Introduction | 8 |
| 2.1. Data versioning and release history | 8 |
| 2.2. Science data processing overview | 9 |
| 2.3. Level 2 products..... | 9 |
| 2.3.1. O3TOT | 9 |
| 2.4. Level 3 product | 10 |
| 3. Science algorithm overview | 11 |
| 3.1. Ancillary data..... | 11 |
| 3.2. O3TOT | 11 |
| 3.2.1. Algorithm overview | 11 |
| 3.2.2. O3TOT usage recommendations | 12 |
| 4. Data Organization and Format | 13 |
| 4.1. Filename..... | 13 |
| 4.2. Format | 13 |
| 4.3. Key variables | 14 |
| 4.3.1. TEMPO Level 2 O3TOT files | 14 |
| References | 16 |

1. Before You Begin

1.1. Purpose of the Document

This document provides a quick guide for Tropospheric Emissions: Monitoring of Pollution (TEMPO) Total Ozone (O3TOT) Level 2 (L2) and Level 3 (L3) products.

This user guide provides the following:

- 1) Algorithm overview
- 2) File descriptions
- 3) Key variables
- 4) Updates from the previous version (Revision history)
- 5) Known issues for the TEMPO O3TOT product

1.2. Acronyms, Abbreviations, and Definitions

| | |
|-------|---|
| AI | Aerosol Index |
| ASDC | Atmospheric Science Data Center |
| ATBD | Algorithm Theoretical Basis Document |
| BUV | Backscatter Ultraviolet |
| CCD | Charge-Coupled Device |
| CF | Cloud Fraction |
| CLDO4 | TEMPO cloud algorithm |
| DU | Dobson Unit |
| FOV | Field of View |
| FWHM | Full Width at Half Maximum |
| HCHO | Formaldehyde |
| INR | Instrument Navigation and Registration |
| L1 | Level 1 |
| L2 | Level 2 |
| L3 | Level 3 |
| LUT | Look-Up Table |
| NASA | National Aeronautics and Space Administration |
| NM | Nadir-Mapper |
| NO2 | Nitrogen Dioxide |
| NPP | National Polar-orbiting Partnership |
| OE | Optimal Estimation |
| OMI | Ozone Monitoring Instrument |

| | |
|--------|---|
| OMPS | Ozone Mapping and Profiler Suite |
| OMTO3 | OMI Level-2 Total Column Ozone Data Product |
| O3TOT | TEMPO total ozone algorithm |
| O3PROF | TEMPO ozone profile algorithm |
| TCO | Total Column Ozone |
| TEMPO | Tropospheric Emissions: Monitoring of Pollution |
| TOA | Top of the Atmosphere |
| TOMS | Total Ozone Mapping Spectrometer |
| QF | Quality Flag |
| SAO | Smithsonian Astrophysical Observatory |
| SZA | Solar Zenith Angle |
| SDPC | Science Data Processing Center |
| TEMPO | Tropospheric Emissions: Monitoring of Pollution |
| TOMS | Total Ozone Mapping Spectrometer |
| UV | Ultraviolet |
| V01 | Version 1 |
| V02 | Version 2 |
| V03 | Version 3 |
| V04 | Version 4 |
| VIS | Visible |
| VZA | Viewing Zenith Angle |

2. Introduction

The Tropospheric Emissions: Monitoring of Pollution (TEMPO) is a geostationary satellite mission to monitor air quality over North America, from Mexico City to the Canadian oil sands and from the Atlantic to the Pacific. It has hourly temporal resolution during sunlight times of the day and high spatial resolution ($2 \text{ km} \times 4.75 \text{ km}$ at the center of the field of regard) (Zoogman et al., 2017).

The TEMPO products are generated by the Science Data Processing Center (SDPC) at the Smithsonian Astrophysical Observatory (SAO). After being generated at SAO, all products are sent to the NASA Atmospheric Science Data Center (ASDC) for public distribution. In rare cases, Level 2 (L2) and Level 3 (L3) product files may be missing even if corresponding Level 1b (L1b) files exist due to failed Instrument Navigation and Registration (INR) processing, which results in a lack of geolocation information.

This User Guide document includes a brief description of the V04 TEMPO L2 and L3 total ozone (O3TOT) products and a quick guide for the V04 TEMPO L2 and L3 O3TOT products, released on September 17, 2025. Note that the user guides for Level 2 and Level 3 ozone profile (O3PROF), trace gas (NO_2 and HCHO), and cloud (CLDO4) products are provided separately.

The SAO team welcomes and relies on feedback from data users to make improvements to TEMPO products.

2.1. Data versioning and release history

Updates to the SDPC operational pipeline result in periodic new data releases. When this occurs, a new collection is created that includes all data products. The collection version number can be found in the data filename and has the format VXX, where XX denotes the collection version. Since first light on August 2, 2023, the SAO science team has been working to improve the processing pipeline and science products with assistance from the TEMPO validation team. On February 5, 2024, a limited set of preliminary unvalidated TEMPO Version 1 (V01) products were released for users to become familiar with the file format and content. On February 26, 2024, the SDPC pipeline was updated to produce Version 2 (V02) products for the public release of the V02 L1b (irradiance and radiance) products. On May 20, 2024, the SDPC was updated to produce Version 3 (V03) L1, L2, and L3 products. V03 products constitute the first public release of TEMPO L2 and L3 products, excluding the TEMPO O3PROF product. As of December 9, 2024, these are declared at the Provisional maturity level. The TEMPO validation document defines the Provisional level as “the performance has been demonstrated through a large, but still (seasonally or otherwise) limited number of independent measurements. The analysis is sufficient for limited qualitative determinations of product fitness-for-purpose, and the product is potentially ready for testing by operational users and may be suitable for scientific publication.” (TEMPO Validation Team, 2023). The provisional maturity level was declared

after validation team approval. Users should consult the relevant data landing pages for current product maturity. On September 17, 2025, the SDPC was updated to produce Version 4 (V04) L1, L2, and L3 products.

2.2. Science data processing overview

The SDPC generates TEMPO products in the following order: (1) L1b, (2) L2 CLDO4, (3) L2 O3TOT, and (4) L3 products. This is because L2 CLDO4 requires L1b as input, L2 products require both L1b and L2 CLDO4 as inputs, and L3 products require L2 products as inputs. Table 2.1.1 lists the L2 and L3 products described in this document. More details for each product can be found in the corresponding ATBD and algorithm description document.

Table 2.1.1. TEMPO L2 and L3 products that are described in this document.

| Product | Level | Description | Spatial Resolution and Coverage | Validation maturity level |
|---------|-------|--|---|---------------------------|
| O3TOT | L2 | Total column ozone (TCO) and UV aerosol index (AI) | Native TEMPO ground pixel, one granule | Provisional |
| O3TOT | L3 | TCO and UV AI on regular grid | All Level 2 granules merged for a single scan, on a $0.02^\circ \times 0.02^\circ$ regular grid | Provisional |

2.3. Level 2 products

2.3.1. O3TOT

The TEMPO O3TOT algorithm is composed of two parts: (1) The forward model calculation for simulating top of atmosphere (TOA) radiance and Jacobians of radiances, (2) The inverse model calculation for deriving the TCO from measured radiance. The TEMPO O3TOT algorithm is based on the OMT03 V8.5 algorithm (called the Total Ozone Mapping Spectrometer (TOMS) V8.5). The detailed description of the algorithm is in the OMI ATBD for ozone products (<https://eospsso.gsfc.nasa.gov/sites/default/files/atbd/ATBD-OMI-02.pdf>, chapter 2). For adapting to the TEMPO, four parts have been updated: (1) Expand viewing geometry to accommodate TEMPO's larger observing viewing zenith angle (VZA), (2) an updated Look-Up Table (LUT) using the TEMPO's on-orbit measured slit functions and BDM (Daumont et al., 1922; Brion et al., 1993; Malicet et al., 1995) ozone absorption cross section, (3) using optical centroid cloud pressure from the TEMPO cloud (CLDO4) product, and (4) performing additional soft calibrations.

2.4. Level 3 product

TEMPO L3 products are generated using an area-weighted gridding program. For each variable of interest, the program maps the corresponding L2 variable onto a $0.02^\circ \times 0.02^\circ$ regular grid over the domain of 155°W – 24.5°W to 17.2°N – 63.55°N . Gridding was performed using data filtered through different solar zenith angles, cloud fraction, and data quality flag criteria as needed by each product.

3. Science algorithm overview

3.1. Ancillary data

TEMPO O3TOT L2 products require ancillary data. The TEMPO O3TOT algorithm mainly uses atmospheric trace gases and meteorological parameters from the TOMS climatology data. They are interpolated onto TEMPO pixels from their original resolutions.

For surface reflectance and effective cloud fraction, TEMPO O3TOT assumes 15% for surface albedo and 80% for cloud albedo to derive effective cloud fraction and directly retrieve surface/cloud albedo if the effective cloud fraction is ≤ 0 or ≥ 1 . Snow and ice can dramatically change surface albedo and significantly influence cloud and trace gas results. Snow and ice fraction is specified through TOMS climatology data. The details of ancillary data are shown in Table 3.1.1.

Table 3.1.1. Ancillary data for TEMPO L2 O3TOT algorithm.

| Input | Source |
|-------------------------|---|
| Cloud-top pressure | TEMPO CLDO4 product Defaults to cloud pressure climatology (OMI-derived) if cloud retrieval is unavailable. |
| Cloud fraction | Retrieved based on the MLER. If $f_c \leq 0$ or ≥ 1 , it is set to 0 or 1, R_s or R_{cl} is retrieved instead. |
| Ozone profiles | TOMS V8 climatology (total ozone dependent, monthly / 10° zonally mean) |
| Temperature profiles | TOMS V8 climatology (monthly / 10° zonally mean) |
| Surface albedo | 15% or directly retrieved as R_s if $f_c = 0$ |
| Snow/ice fraction | Climatology at $1^\circ \times 1^\circ$ from the TOMS V8 |
| Terrain height pressure | Climatology at $1/3^\circ \times 1/3^\circ$ from the TOMS V8 |
| Aerosols | Not explicitly treated, but an aerosol correction is included |

3.2. O3TOT

3.2.1. Algorithm overview

The TEMPO L2 O3TOT algorithm is based on the OMT03 V8.5 algorithm (called TOMS V8.5) adapted for TEMPO. This algorithm has been described in detail in the OMI ATBD for ozone products (<https://eospsso.gsfc.nasa.gov/sites/default/files/atbd/ATBD-OMI-02.pdf>, chapter 2).

There are several changes to adapt this algorithm for TEMPO L2 O3TOT. The first way to adapt to TEMPO is to update the radiance Look-Up Table (LUT). The viewing zenith angles in

LUT were expanded to accommodate TEMPO's larger viewing zenith angle (VZA). An additional improvement in the V04 TEMPO O3TOT product involves updating the radiance LUT by incorporating the on-orbit measured TEMPO slit function, which more accurately characterizes the instrument's spectral response. This updated LUT is applied in combination with the BDM ozone absorption cross sections. Additionally, the optical centroid cloud pressure from the TEMPO cloud (CLDO4) product is utilized. The last way to adapt to TEMPO is performing the additional soft calibration.

The basic algorithm uses 2 wavelengths (317.5 and 331.2 nm under most conditions, and 331.2 and 360 nm for high ozone and high solar zenith angle conditions). The longer of the two wavelengths is used to derive effective cloud fraction (f_c) based on the Mixed Lambert Equivalent Reflectivity (MLER) model that was developed to model the effect of clouds on Rayleigh scattering. This algorithm also calculates the absorbing aerosol index (AI) from the radiance residuals at 360 nm.

3.2.2. O3TOT usage recommendations

The `quality_flag` variable in the product group of the L2 file provides the quality of the product. Retrieval of the highest quality has a `quality_flag` equal to "0", a solar zenith angle (SZA) less than 80 degrees, a viewing zenith angle (VZA) less than 80 degrees, and a cloud fraction less than 0.5.

Suppose a data user aims to perform an analysis with a larger volume of data, even if it is less accurate. In that case, the data with `quality_flag` values of "1" (Glint contamination), "2" (SZA greater than 84 degrees), or "5" (SOI greater than 4 sigma) can be used.

To ensure reasonable data quality, it is recommended to additionally filter for SZA and VZA less than 80 degrees, and an effective cloud fraction below 0.5.

Lastly, if a data user wants to perform an analysis without considering the data quality, the user can use a `quality_flag` that is smaller than "4096" (Warning input Radiance).

In the case of the TEMPO L3 O3TOT product. Because the meaning of the `quality_flag` could change after averaging pixels, we can't provide the `quality_flag` variable in the TEMPO L3 O3TOT product. Therefore, the TEMPO L3 O3TOT product applied data filtering as described above before producing the TEMPO L3 O3TOT product.

4. Data Organization and Format

TEMPO data products are organized in terms of the scan number (e.g., S004) and granule number (e.g., G01) for each day. Each scan is a sweep from the East to the West, and nominally takes one hour to complete. The data collected during each scan is divided into different granules to keep the data volume for each file manageable. Each granule contains data collected within a few minutes.

4.1. Filename

Table 4.1.1 shows the TEMPO Level 2 and 3 file naming convention. The Level 2 O3TOT filenames have a format of

TEMPO_{GAS}_L2_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}G{YY}.nc,
with YYYY (year), MM (month), DD (day), HH (hour), NN (minute), and SS (second) denoting the GMT time stamp at the beginning of the time period covered by the file, and XXX and YY denoting the scan and granule numbers, respectively. The segment GAS represents the product type: O3TOT. (Note that the curly brackets are added here for display purposes; they don't exist in actual filenames.)

The TEMPO Level 3 filenames have a format of

TEMPO_{GAS}_L3_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}.nc,

Again, with YYYY (year), MM (month), DD (day), HH (hour), NN (minute), and SS (second) denoting the GMT time stamp at the beginning of the time period covered by the file and _S{XXX} the scan number.

Table 4.1.1 TEMPO Level 2 and 3 file naming convention.

| Product | Filename format |
|------------------|--|
| O3TOT Level 2 | TEMPO_O3TOT_L2_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}G{YY}.nc |
| O3TOT Level 3 | TEMPO_O3TOT_L3_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}.nc |

4.2. Format

TEMPO products are in the self-explanatory NetCDF4 format. The content of a NetCDF file can be reviewed by a command “ncdump -h {FILENAME}” on Linux systems or explored on a graphical interface using the software “Panoply,” distributed by NASA Goddard Institute for

Space Studies (<https://www.giss.nasa.gov/tools/panoply/>). NetCDF files can be read using FORTRAN, C, Python, Matlib, IDL, etc.

Each file contains several groups (named “diagnostic”, “geolocation”, “product”, “qa_statistics”, and “support_data”) and global attributes. Each group contains a number of variables. Each variable is described by its attributes.

Dimensions used for the variables are defined in each netcdf file. Typically, the xtrack = 2048 dimension is fixed along the North-South direction. The first and last few pixels in the North-South direction contain fill values as they are not suitable for retrieval. The granule builds in the East-West scanning direction, and contains data collected within a few minutes. The corner = 4 dimension is used for the 4 corners of each ground pixel.

4.3. Key variables

Key variables for TEMPO L2 and L3 files in this release are provided below. For the full list of variables, please use NetCDF manipulation software.

4.3.1. TEMPO Level 2 O3TOT files

Table 4.3.1. TEMPO Level 2 O3TOT key variables

| Group | Name | Format (dimension) | Unit | Description |
|---------|------------------|--------------------------------|------|---|
| product | column_amount_o3 | float (mirror_step, xtrack) | DU | Total ozone column |
| product | fc | float (mirror_step, xtrack) | none | Effective cloud fraction (mixed LER model) |
| product | uv_aerosol_index | float (mirror_step, xtrack) | none | UV aerosol index |

The L2 O3TOT file also contains multiple useful variables to check the quality of the total ozone retrieval. The following variables are included in O3TOT L2 file:

- quality_flag in product group reports the detailed quality of the retrieval.
 - Bits 0 to 3 together contain several output error flags (Note in the case of multiple error flags, only the highest error flag is reported):
 - 0: Good sample
 - 1: Glint contamination (corrected)
 - 2: SZA > 84°
 - 3: 360 residual > threshold
 - 4: Residual at unused ozone wavelength > 4 sigma
 - 5: SOI > 4 sigma (SO₂ present)

- 6: Non-convergence
 - 7: Abs (residual) > 16.0 (fatal)
 - 8: Row anomaly error (same as bit 6 in this field)
 - Bits 4 to 5 are reserved for future use (currently set to 0).
 - Bit 7 is set to 0 when TEMPO CLDO4 cloud pressure is used and set to 1 when climatological cloud pressure is used.
 - Bits 8 to 15 are flags that are set to 0 for FALSE (good value), or 1 for TRUE (bad value)
 - Bit 8: Geolocation error (anomalous FOV Earth location)
 - Bit 9: SZA > 88°
 - Bit 10: Missing input radiance
 - Bit 11: Error input radiance
 - Bit 12: Warning input Radiance
 - Bit 13: Missing input irradiance
 - Bit 14: Error input irradiance
 - Bit 15: Warning input irradiance
- algorithm_flags in support_data group reports the overall quality of the retrieval.
 - The algorithm flag associated with the ground pixel:
 - 0: Skipped
 - 1: Standard
 - 2: Adjusted for profile shape
 - 3: Based on C-pair (331 and 360 nm)
 - 10: Snow/Ice

References

- Bass, A. M., & Paur, R. J. (1985). The ultraviolet cross-sections of ozone: I. The measurements. In *Atmospheric Ozone: Proceedings of the Quadrennial Ozone Symposium held in Halkidiki, Greece 3–7 September 1984* (pp. 606-610). Dordrecht: Springer Netherlands.
https://doi.org/10.1007/978-94-009-5313-0_120
- Brion, J., Chakir, A., Daumont, D., Malicet, J., & Parisse, C. (1993). High-resolution laboratory absorption cross section of O₃. Temperature effect. *Chemical physics letters*, 213(5-6), 610-612. [https://doi.org/10.1016/0009-2614\(93\)89169-I](https://doi.org/10.1016/0009-2614(93)89169-I)
- Bhartia, P. K., & Wellemeyer, C. W. (2002). *TOMS-V8 total O₃ algorithm. OMI Algorithm Theoretical Basis Document Volume II, NASA Goddard Space Flight Center Tech. Doc. ATBD-OMI-02*, 15–31, <https://eosps.gsfc.nasa.gov/sites/default/files/atbd/ATBD-OMI-02.pdf>.
- Chong, H., Liu, X., Houck, J., Flittner, D. E., Carr, J., Hou, W., Suleiman, R. M., & Chance, K. (2024). TEMPO Level 1 Data Product: User Guide.
- Chong, H., Liu, X., Houck, J., Flittner, D. E., Carr, J., Hou, W., Davis, J. E., Suleiman, R. M., Chance, K., Mishra, N., Chan Miller, C., González Abad, G., Baker, B., Lasnik, J., Nicks, D., Bak, J., Nowlan, C. R., Wang, H., Park, J., O’Sullivan, E., Fitzmaurice, J., & Carpenter, L. (2024). Algorithm theoretical basis document for the TEMPO Level 0-1 processor.
- Daumont, D., Brion, J., Charbonnier, J., & Malicet, J. (1992). Ozone UV spectroscopy I: Absorption cross-sections at room temperature. *Journal of Atmospheric Chemistry*, 15, 145-155. <https://doi.org/10.1007/BF00053756>
- Dave, J. V., & Mateer, C. L. (1967). A preliminary study on the possibility of estimating total atmospheric ozone from satellite measurements. *Journal of the Atmospheric Sciences*, 24(4), 414-427. [https://doi.org/10.1175/1520-0469\(1967\)024<0414:APSOTP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1967)024<0414:APSOTP>2.0.CO;2)

- Malicet, J., Daumont, D., Charbonnier, J., Parisse, C., Chakir, A., & Brion, J. (1995). Ozone UV spectroscopy. II. Absorption cross-sections and temperature dependence. *Journal of atmospheric chemistry*, 21, 263-273. <https://doi.org/10.1007/BF00696758>
- Park, J., Liu, X., Bak, J., Houck, J., Chance, K., Suleiman, R. M., Davis, J. E., Chong, H., Hou, W., Flittner, D. E., Carr, J., O'Sullivan, E., González Abad, G., Knowland, K. E., Chan Miller, C., Nowlan, C. R., Wang, H., Fitzmaurice, J., Carpenter, L., Spurr, R., & Newchurch, M. J. (2024). Algorithm theoretical basis document for the TEMPO Ozone Profile Retrieval Algorithm.
- Park, J., Liu, X., Houch, J., Haffner, D., & Chnace, K. (2024). Algorithm Description for the TEMPO Total Ozone Retrieval Algorithm.
- TEMPO Validation Team (2023). Tropospheric Emissions: Monitoring of Pollution (TEMPO) – Level 2 Science Data Product Validation Plan, SAO-DRD-11.
- Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Liu, X., Park, J., Petropavlovskikh, I., Hanisco, T.F., Hanisco, T., Szykman, J., Valin, L., Baumann, E., Cede, A., Tiefengraber, M., Gebetsberger, M., Uesato, I., Zheng, X., Ahn, S., Chang, L., Lee, W., Kim, J., Lee, H., Baek, K., Redondas, A., Fujiwara, M., Wang, T., Grutter, M., Houck, J., Haffner, D., & Lee, S. (2025). Geostationary Satellites Total Ozone Observations: First Results on Ground-based Networks Validation Efforts for TEMPO and GEMS. *Geophysical Research Letters*, 52, e2025GL114768. <https://doi.org/10.1029/2025GL114768>.
- Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A., Hilton, B. B., Nicks, D. K., Newchurch, M. J., Carr, J. L. Janz, S. J., Andraschko, M. R., Arola, A., Baker, B. D., Canova, B. P., Chan Miller, C., Cohen, R. C., Davis, J. E., Dussault, M. E., Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter, M., Herman, J. R., Houck, J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C., Lindfors, A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R., O'Sullivan, E. J., Palmer, P. I., Pierce, R. B., Pippin, M. R., Saiz-Lopez, A., Spurr, R. J. D., Szykman, J. J., Torres, O., Veefkind, J. P., Veihelmann, B., Wang, H., Wang, J., & Chance, K. (2017). Tropospheric emissions:

Monitoring of pollution (TEMPO). *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 17-39. <https://doi.org/10.1016/j.jqsrt.2016.05.008>