



CENTER FOR
ASTROPHYSICS
HARVARD & SMITHSONIAN

VERSION: 2.2
RELEASE DATE: February 23, 2026



**TROPOSPHERIC EMISSIONS:
MONITORING OF POLLUTION (TEMPO)
PROJECT**

**Total Ozone Level 2 and 3 Data products:
User Guide**

February 23, 2026

AUTHORS

Junsung Park (Center for Astrophysics | Harvard & Smithsonian)

Xiong Liu (Center for Astrophysics | Harvard & Smithsonian)

John Houck (Center for Astrophysics | Harvard & Smithsonian)

Dave Haffner (NASA Goddard Space Flight Center, Science Systems and Applications Inc.)

Kelly Chance (Center for Astrophysics | Harvard & Smithsonian)

CONTRIBUTIONS:

TEMPO Science Team, Center for Astrophysics | Harvard & Smithsonian

TEMPO Ground System Team, Center for Astrophysics | Harvard & Smithsonian

NASA Langley Research Center

NASA Goddard Space Flight Center

REVISION HISTORY

Version	Item	Description	Release date
1.0	All	Initial version	May 20, 2024
1.1	Known data product issues	Update known data product issues	December 9, 2024
	Section 2	Update validation status to provisional	
	Section 3.2.2	Add a Section for data versioning and release history	
	Section 4.3	Update quality flag explanation	
2.0	Known data product issues	Update known data product issues	September 17, 2025
	Section 2	Update Algorithm overview	
	Section 3	Update O3TOT usage recommendations	
2.1	Document version	Update the document version	January 6 2026
	Revision history	Update revision history	
	Section 3.2.1	Update the wavelength informations	
	Section 3.2.2	Update the usage recommendations	
	Reference	Update the references	
2.2	Known data product issues	Update known data product issues	February 23, 2026
	Section 3.2.1	Update the OMI ATBD reference link and correct the UVAI retrieval explanation	
	All	Correct the resolution of L3 product	

PRODUCT to SCIENCE DATA PROCESSING CENTER PIPELINE VERSION

Version Designation	Science Data Processing Center Pipeline Version	Release
V03	4.4	May 20, 2024; first public release of TEMPO Level 2 and Level 3 products
V04	4.7	September 17, 2025

KNOWN DATA PRODUCT ISSUES

Total Ozone (O3TOT)

- (1) TEMPO V03 O3TOT Total column ozone (TCO) shows latitude-dependent biases compared to OMI and OMPS TCO, varying from ~5% at lower latitudes to ~-2% at high latitudes. This latitudinal dependence does not vary much with season. This latitudinal dependence is confirmed from validation using ground-based Pandora, Dobson, and Brewer observations (Zhao et al., 2025). Preliminary testing indicates that this latitudinal dependence results mainly from the spectral dependence of radiometric bias in the Sun-normalized radiances. In addition, V03 O3TOT UV aerosol index shows negative biases of ~-2, primarily from the large overestimation in the normalized radiance of V03 L1b data.
 - a. This latitude dependence is caused by an issue in the TEMPO L1B data, which exhibits an additional bump feature in the irradiance spectrum between 300 nm and 330 nm after a wavelength-independent reduction of normalized radiance by ~13.4% in the UV band of V04 L1b data. To correct this, we implemented soft calibration at the wavelengths used in the ozone retrieval (312.61 nm and 317.62 nm). Additionally, we performed soft calibration to the wavelength (360.15 nm) used in the reflectance and aerosol index calculation.
 - b. After applying these soft calibrations after the L1b update, the latitude dependence has been mitigated in the O3TOT V04 data product. Preliminary evaluation indicates that the TCO from the TEMPO O3TOT algorithm shows excellent agreement, with a relative difference within $\pm 1\%$, with TCO from NPP OMPS satellite observations.
 - c. The peak of the histogram of UV aerosol index is improved from ~-2.5 to ~0, mainly from the L1b update and also slightly from additional soft calibrations.
- (2) The TEMPO V04 O3TOT TCO product shows improvements in latitude-dependent biases compared to V03 O3TOT TCO, while still showing a slight negative bias.
 - a. An evaluation of TCO from the publicly released TEMPO V04 O3TOT product against Pandora observations at 53 Pandonia Global Network (PGN) sites across North America, covering the period from September 17 to December 31, 2025, demonstrates a correlation coefficient of 0.98, a Mean Bias (MB) of -2.13%, and an RMSE of 2.74%.
 - b. TEMPO V04 O3TOT TCO shows a consistent negative bias compared to OMPS and TROPOMI TCO observations. A larger negative bias in high-latitude regions is likely associated with large solar and viewing zenith angles (SZA and VZA). Future algorithm updates will address this issue by optimizing the C-pair selection threshold and the profile-shape switching threshold, which are expected to reduce systematic biases and improve performance at large viewing angles.

- (3) The quality_flag has an issue (including unnecessary flags for the TEMPO observation) in the V03 TEMPO O3TOT product.
 - a. This issue has been updated from the V04 TEMPO O3TOT product. Therefore, users can confidently use good samples with quality_flag set to 0 (with $SZA \leq 80$, $VZA \leq 80$, and $CF \leq 0.5$).
- (4) TEMPO TCO also shows clear discontinuity when the algorithm switches from using the main pair of wavelengths (~317 and 331 nm) to a different wavelength pair (331 and 360 nm) at high solar zenith angle or viewing zenith angle, resulting in lower TCO values.
 - a. To address this, the thresholds for wavelength switch and profile shape change will be updated to reflect TEMPO characteristics. These thresholds will determine the appropriate values using a sufficiently large set of TEMPO O3TOT V04 product and incorporate them in the next product version.

Contents

AUTHORS	2
REVISION HISTORY	3
PRODUCT to SCIENCE DATA PROCESSING CENTER PIPELINE VERSION	3
KNOWN DATA PRODUCT ISSUES	4
Total Ozone (O3TOT)	4
1. Before You Begin.....	7
1.1. Purpose of the Document.....	7
1.2. Acronyms, Abbreviations, and Definitions	7
2. Introduction	9
2.1. Data versioning and release history	9
2.2. Science data processing overview	10
2.3. Level 2 products.....	10
2.3.1. O3TOT	10
2.4. Level 3 product	11
3. Science algorithm overview	12
3.1. Ancillary data.....	12
3.2. O3TOT	12
3.2.1. Algorithm overview	12
3.2.2. O3TOT usage recommendations	13
4. Data Organization and Format	14
4.1. Filename.....	14
4.2. Format.....	14
4.3. Key variables	15
4.3.1. TEMPO Level 2 O3TOT files	15
References	17

1. Before You Begin

1.1. Purpose of the Document

This document provides a quick guide for Tropospheric Emissions: Monitoring of Pollution (TEMPO) Total Ozone (O3TOT) Level 2 (L2) and Level 3 (L3) products.

This user guide provides the following:

- 1) Algorithm overview
- 2) File descriptions
- 3) Key variables
- 4) Updates from the previous version (Revision history)
- 5) Known issues for the TEMPO O3TOT product

1.2. Acronyms, Abbreviations, and Definitions

AI	Aerosol Index
ASDC	Atmospheric Science Data Center
ATBD	Algorithm Theoretical Basis Document
BUV	Backscatter Ultraviolet
CCD	Charge-Coupled Device
CF	Cloud Fraction
CLDO4	TEMPO cloud algorithm
DU	Dobson Unit
FOV	Field of View
FWHM	Full Width at Half Maximum
HCHO	Formaldehyde
INR	Instrument Navigation and Registration
L1	Level 1
L2	Level 2
L3	Level 3
LUT	Look-Up Table
NASA	National Aeronautics and Space Administration
NM	Nadir-Mapper
NO2	Nitrogen Dioxide
NPP	National Polar-orbiting Partnership

OE	Optimal Estimation
OMI	Ozone Monitoring Instrument
OMPS	Ozone Mapping and Profiler Suite
OMTO3	OMI Level-2 Total Column Ozone Data Product
O3TOT	TEMPO total ozone algorithm
O3PROF	TEMPO ozone profile algorithm
TCO	Total Column Ozone
TEMPO	Tropospheric Emissions: Monitoring of Pollution
TOA	Top of the Atmosphere
TOMS	Total Ozone Mapping Spectrometer
QF	Quality Flag
SAO	Smithsonian Astrophysical Observatory
SZA	Solar Zenith Angle
SDPC	Science Data Processing Center
TEMPO	Tropospheric Emissions: Monitoring of Pollution
TOMS	Total Ozone Mapping Spectrometer
UV	Ultraviolet
V01	Version 1
V02	Version 2
V03	Version 3
V04	Version 4
VIS	Visible
VZA	Viewing Zenith Angle

2. Introduction

The Tropospheric Emissions: Monitoring of Pollution (TEMPO) is a geostationary satellite mission to monitor air quality over North America, from Mexico City to the Canadian oil sands and from the Atlantic to the Pacific. It has hourly temporal resolution during sunlight times of the day and high spatial resolution ($2 \text{ km} \times 4.75 \text{ km}$ at the center of the field of regard) (Zoogman et al., 2017).

The TEMPO products are generated by the Science Data Processing Center (SDPC) at the Smithsonian Astrophysical Observatory (SAO). After being generated at SAO, all products are sent to the NASA Atmospheric Science Data Center (ASDC) for public distribution. In rare cases, Level 2 (L2) and Level 3 (L3) product files may be missing even if corresponding Level 1b (L1b) files exist due to failed Instrument Navigation and Registration (INR) processing, which results in a lack of geolocation information.

This User Guide document includes a brief description of the V04 TEMPO L2 and L3 total ozone (O3TOT) products and a quick guide for the V04 TEMPO L2 and L3 O3TOT products, released on September 17, 2025. Note that the user guides for Level 2 and Level 3 ozone profile (O3PROF), trace gas (NO₂ and HCHO), and cloud (CLDO4) products are provided separately.

The SAO team welcomes and relies on feedback from data users to make improvements to TEMPO products.

2.1. Data versioning and release history

Updates to the SDPC operational pipeline result in periodic new data releases. When this occurs, a new collection is created that includes all data products. The collection version number can be found in the data filename and has the format VXX, where XX denotes the collection version. Since first light on August 2, 2023, the SAO science team has been working to improve the processing pipeline and science products with assistance from the TEMPO validation team. On February 5, 2024, a limited set of preliminary unvalidated TEMPO Version 1 (V01) products were released for users to become familiar with the file format and content. On February 26, 2024, the SDPC pipeline was updated to produce Version 2 (V02) products for the public release of the V02 L1b (irradiance and radiance) products. On May 20, 2024, the SDPC was updated to produce Version 3 (V03) L1, L2, and L3 products. V03 products constitute the first public release of TEMPO L2 and L3 products, excluding the TEMPO O3PROF product. As of December 9, 2024, these are declared at the Provisional maturity level. The TEMPO validation document defines the Provisional level as “the performance has been demonstrated through a large, but still (seasonally or otherwise) limited number of independent measurements. The analysis is sufficient for limited qualitative determinations of product fitness-for-purpose, and the product is potentially ready for testing by operational users and may be suitable for scientific publication.” (TEMPO Validation Team, 2023). The provisional maturity level was declared

after validation team approval. Users should consult the relevant data landing pages for current product maturity. On September 17, 2025, the SDPC was updated to produce Version 4 (V04) L1, L2, and L3 products.

2.2. Science data processing overview

The SDPC generates TEMPO products in the following order: (1) L1b, (2) L2 CLDO4, (3) L2 O3TOT, and (4) L3 products. This is because L2 CLDO4 requires L1b as input, L2 products require both L1b and L2 CLDO4 as inputs, and L3 products require L2 products as inputs. Table 2.1.1 lists the L2 and L3 products described in this document. More details for each product can be found in the corresponding ATBD and algorithm description document.

Table 2.1.1. TEMPO L2 and L3 products that are described in this document.

Product	Level	Description	Spatial Resolution and Coverage	Validation maturity level
O3TOT	L2	Total column ozone (TCO) and UV aerosol index (AI)	Native TEMPO ground pixel, one granule	Provisional
O3TOT	L3	TCO and UV AI on regular grid	All Level 2 granules merged for a single scan, on a 0.02° (longitude) × 0.04° (latitude) regular grid	Provisional

2.3. Level 2 products

2.3.1. O3TOT

The TEMPO O3TOT algorithm is composed of two parts: (1) The forward model calculation for simulating top of atmosphere (TOA) radiance and Jacobians of radiances, (2) The inverse model calculation for deriving the TCO from measured radiance. The TEMPO O3TOT algorithm is based on the OMT03 V8.5 algorithm (called the Total Ozone Mapping Spectrometer (TOMS) V8.5). The detailed description of the algorithm is in the OMI ATBD for ozone products (<https://ozoneaq.gsfc.nasa.gov/content/public/OMI/ATBDs/ATBD-OMI-02.pdf>, chapter 2). For adapting to the TEMPO, four parts have been updated: (1) Expand viewing geometry to accommodate TEMPO's larger observing viewing zenith angle (VZA), (2) an updated Look-Up Table (LUT) using the TEMPO's on-orbit measured slit functions and BDM (Daumont et al., 1922; Brion et al., 1993; Malicet et al., 1995) ozone absorption cross section, (3) using optical centroid cloud pressure from the TEMPO cloud (CLDO4) product, and (4) performing additional soft calibrations.

2.4. Level 3 product

TEMPO L3 products are generated using an area-weighted gridding program. For each variable of interest, the program maps the corresponding L2 variable onto a $0.02^\circ \times 0.04^\circ$ regular grid over the domain of 155°W – 24.5°W to 17.2°N – 63.55°N . Gridding was performed using data filtered through different solar zenith angles, cloud fraction, and data quality flag criteria as needed by each product.

3. Science algorithm overview

3.1. Ancillary data

TEMPO O3TOT L2 products require ancillary data. The TEMPO O3TOT algorithm mainly uses atmospheric trace gases and meteorological parameters from the TOMS climatology data. They are interpolated onto TEMPO pixels from their original resolutions.

For surface reflectance and effective cloud fraction, TEMPO O3TOT assumes 15% for surface albedo and 80% for cloud albedo to derive effective cloud fraction and directly retrieve surface/cloud albedo if the effective cloud fraction is ≤ 0 or ≥ 1 . Snow and ice can dramatically change surface albedo and significantly influence cloud and trace gas results. Snow and ice fraction is specified through TOMS climatology data. The details of ancillary data are shown in Table 3.1.1.

Table 3.1.1. Ancillary data for TEMPO L2 O3TOT algorithm.

Input	Source
Cloud-top pressure	TEMPO CLDO4 product Defaults to cloud pressure climatology (OMI-derived) if cloud retrieval is unavailable.
Cloud fraction	Retrieved based on the MLER. If $f_c \leq 0$ or ≥ 1 , it is set to 0 or 1, R_s or R_{cl} is retrieved instead.
Ozone profiles	TOMS V8 climatology (total ozone dependent, monthly / 10° zonally mean)
Temperature profiles	TOMS V8 climatology (monthly / 10° zonally mean)
Surface albedo	15% or directly retrieved as R_s if $f_c = 0$
Snow/ice fraction	Climatology at 1° × 1° from the TOMS V8
Terrain height pressure	Climatology at 1/3° × 1/3° from the TOMS V8
Aerosols	Not explicitly treated, but an aerosol correction is included

3.2. O3TOT

3.2.1. Algorithm overview

The TEMPO L2 O3TOT algorithm is based on the OMT03 V8.5 algorithm (called TOMS V8.5) adapted for TEMPO. This algorithm has been described in detail in the OMI ATBD for ozone products (<https://ozoneaq.gsfc.nasa.gov/content/public/OMI/ATBDs/ATBD-OMI-02.pdf>, chapter 2).

There are several changes to adapt this algorithm for TEMPO L2 O3TOT. The first way to adapt to TEMPO is to update the radiance Look-Up Table (LUT). The viewing zenith angles in

LUT were expanded to accommodate TEMPO's larger viewing zenith angle (VZA). An additional improvement in the V04 TEMPO O3TOT product involves updating the radiance LUT by incorporating the on-orbit measured TEMPO slit function, which more accurately characterizes the instrument's spectral response. This updated LUT is applied in combination with the BDM ozone absorption cross sections. Additionally, the optical centroid cloud pressure from the TEMPO cloud (CLDO4) product is utilized. The last way to adapt to TEMPO is performing the additional soft calibration.

The basic algorithm uses 2 wavelengths (317.62 and 331.34 nm under most conditions, and 331.34 and 360.15 nm for high ozone and high solar zenith angle conditions). The longer of the two wavelengths is used to derive effective cloud fraction (f_c) based on the Mixed Lambert Equivalent Reflectivity (MLER) model that was developed to model the effect of clouds on Rayleigh scattering. This algorithm, consistent with the TOMS V8 heritage, identifies absorbing aerosols. The absorbing Aerosol Index (AI) is defined as the logarithmic contrast between measured and Rayleigh-calculated radiances at 360.15 and 331.34 nm. Reflectivity derived at 331.34 nm is used to compute the expected radiance at 360.15 nm under a Rayleigh-only atmosphere using precomputed lookup tables. Positive AI values indicate the presence of UV-absorbing aerosols.

3.2.2. O3TOT usage recommendations

The `quality_flag` variable in the product group of the L2 file provides the quality of the product. Retrieval of the highest quality has a `quality_flag` equal to "0", a solar zenith angle (SZA) less than 80 degrees, a viewing zenith angle (VZA) less than 80 degrees, and a cloud fraction less than 0.5.

Suppose a data user aims to perform an analysis with a larger volume of data, even if it is less accurate. In that case, the data with `quality_flag` values of "1" (Glint contamination), "2" (SZA greater than 84 degrees), or "5" (SOI greater than 4 sigma) can be used.

To ensure reasonable data quality, it is recommended to additionally filter for SZA and VZA less than 80 degrees, and an effective cloud fraction below 0.5.

Lastly, if a data user wants to perform an analysis without considering the data quality, the user can use a `quality_flag` that is smaller than "1024" (Missing, Error, or Warning input Radiance/Irradiance).

In the case of the TEMPO L3 O3TOT product. Because the meaning of the `quality_flag` could change after averaging pixels, we can't provide the `quality_flag` variable in the TEMPO L3 O3TOT product. Therefore, the TEMPO L3 O3TOT product applied data filtering as described above before producing the TEMPO L3 O3TOT product. Additional filtering using cloud fraction could be applied for better quality.

4. Data Organization and Format

TEMPO data products are organized in terms of the scan number (e.g., S004) and granule number (e.g., G01) for each day. Each scan is a sweep from the East to the West, and nominally takes one hour to complete. The data collected during each scan is divided into different granules to keep the data volume for each file manageable. Each granule contains data collected within a few minutes.

4.1. Filename

Table 4.1.1 shows the TEMPO Level 2 and 3 file naming convention. The Level 2 O3TOT filenames have a format of

TEMPO_{GAS}_L2_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}G{YY}.nc,
with YYYY (year), MM (month), DD (day), HH (hour), NN (minute), and SS (second) denoting the GMT time stamp at the beginning of the time period covered by the file, and XXX and YY denoting the scan and granule numbers, respectively. The segment GAS represents the product type: O3TOT. (Note that the curly brackets are added here for display purposes; they don't exist in actual filenames.)

The TEMPO Level 3 filenames have a format of

TEMPO_{GAS}_L3_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}.nc,
Again, with YYYY (year), MM (month), DD (day), HH (hour), NN (minute), and SS (second) denoting the GMT time stamp at the beginning of the time period covered by the file and _S{XXX} the scan number.

Table 4.1.1 TEMPO Level 2 and 3 file naming convention.

Product	Filename format
O3TOT Level 2	TEMPO_O3TOT_L2_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}G{YY}.nc
O3TOT Level 3	TEMPO_O3TOT_L3_V04_{YYYY}{MM}{DD}T{HH}{NN}{SS}Z_S{XXX}.nc

4.2. Format

TEMPO products are in the self-explanatory NetCDF4 format. The content of a NetCDF file can be reviewed by a command “ncdump -h {FILENAME}” on Linux systems or explored on a graphical interface using the software “Panoply,” distributed by NASA Goddard Institute for

Space Studies (<https://www.giss.nasa.gov/tools/panoply/>). NetCDF files can be read using FORTRAN, C, Python, Matlab, IDL, etc.

Each file contains several groups (named “diagnostic”, “geolocation”, “product”, “qa_statistics”, and “support_data”) and global attributes. Each group contains a number of variables. Each variable is described by its attributes.

Dimensions used for the variables are defined in each netcdf file. Typically, the xtrack = 2048 dimension is fixed along the North-South direction. The first and last few pixels in the North-South direction contain fill values as they are not suitable for retrieval. The granule builds in the East-West scanning direction, and contains data collected within a few minutes. The corner = 4 dimension is used for the 4 corners of each ground pixel.

4.3. Key variables

Key variables for TEMPO L2 and L3 files in this release are provided below. For the full list of variables, please use NetCDF manipulation software.

4.3.1. TEMPO Level 2 O3TOT files

Table 4.3.1. TEMPO Level 2 O3TOT key variables

Group	Name	Format (dimension)	Unit	Description
product	column_amount_o3	float (mirror_step, xtrack)	DU	Total ozone column
product	fc	float (mirror_step, xtrack)	none	Effective cloud fraction (mixed LER model)
product	uv_aerosol_index	float (mirror_step, xtrack)	none	UV aerosol index

The L2 O3TOT file also contains multiple useful variables to check the quality of the total ozone retrieval. The following variables are included in O3TOT L2 file:

- quality_flag in product group reports the detailed quality of the retrieval.
 - Bits 0 to 3 together contain several output error flags (Note in the case of multiple error flags, only the highest error flag is reported):
 - 0: Good sample
 - 1: Glint contamination (corrected)
 - 2: SZA > 84°
 - 3: 360 residual > threshold
 - 4: Residual at unused ozone wavelength > 4 sigma
 - 5: SOI > 4 sigma (SO₂ present)

- 6: Non-convergence
 - 7: Abs (residual) > 16.0 (fatal)
 - 8: Row anomaly error (same as bit 6 in this field)
 - Bits 4 to 5 are reserved for future use (currently set to 0).
 - Bit 7 is set to 0 when TEMPO CLDO4 cloud pressure is used and set to 1 when climatological cloud pressure is used.
 - Bits 8 to 15 are flags that are set to 0 for FALSE (good value), or 1 for TRUE (bad value)
 - Bit 8: Geolocation error (anomalous FOV Earth location)
 - Bit 9: SZA > 88°
 - Bit 10: Missing input radiance
 - Bit 11: Error input radiance
 - Bit 12: Warning input Radiance
 - Bit 13: Missing input irradiance
 - Bit 14: Error input irradiance
 - Bit 15: Warning input irradiance
- algorithm_flags in support_data group reports the overall quality of the retrieval.
 - The algorithm flag associated with the ground pixel:
 - 0: Skipped
 - 1: Standard
 - 2: Adjusted for profile shape
 - 3: Based on C-pair (331 and 360 nm)
 - 10: Snow/Ice

References

- Bass, A. M., & Paur, R. J. (1985). The ultraviolet cross-sections of ozone: I. The measurements. In *Atmospheric Ozone: Proceedings of the Quadrennial Ozone Symposium held in Halkidiki, Greece 3–7 September 1984* (pp. 606-610). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-009-5313-0_120
- Brion, J., Chakir, A., Daumont, D., Malicet, J., & Parisse, C. (1993). High-resolution laboratory absorption cross section of O₃. Temperature effect. *Chemical physics letters*, 213(5-6), 610-612. [https://doi.org/10.1016/0009-2614\(93\)89169-I](https://doi.org/10.1016/0009-2614(93)89169-I)
- Bhartia, P. K., & Wellemeyer, C. W. (2002). *TOMS-V8 total O₃ algorithm. OMI Algorithm Theoretical Basis Document Volume II, NASA Goddard Space Flight Center Tech. Doc. ATBD-OMI-02*, 15–31. <https://ozoneaq.gsfc.nasa.gov/content/public/OMI/ATBDs/ATBD-OMI-02.pdf>
- Chong, H., Liu, X., Houck, J., Flittner, D. E., Carr, J., Hou, W., Suleiman, R. M., & Chance, K. (2026). TEMPO Level 1 Data Product: User Guide. https://asdc.larc.nasa.gov/documents/tempo/guide/TEMPO_Level-1_user_guide_V2.0.pdf
- Chong, H., Liu, X., Houck, J., Flittner, D. E., Carr, J., Hou, W., Davis, J. E., Suleiman, R. M., Chance, K., Mishra, N., Chan Miller, C., González Abad, G., Baker, B., Lasnik, J., Nicks, D., Bak, J., Nowlan, C. R., Wang, H., Park, J., O’Sullivan, E., Fitzmaurice, J., & Carpenter, L. (2026). Algorithm theoretical basis document for the TEMPO Level 0-1 processor. <https://doi.org/10.1029/2025EA004516>
- Daumont, D., Brion, J., Charbonnier, J., & Malicet, J. (1992). Ozone UV spectroscopy I: Absorption cross-sections at room temperature. *Journal of Atmospheric Chemistry*, 15, 145-155. <https://doi.org/10.1007/BF00053756>
- Dave, J. V., & Mateer, C. L. (1967). A preliminary study on the possibility of estimating total atmospheric ozone from satellite measurements. *Journal of the Atmospheric Sciences*, 24(4), 414-427. [https://doi.org/10.1175/1520-0469\(1967\)024<0414:APSOTP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1967)024<0414:APSOTP>2.0.CO;2)

- Malicet, J., Daumont, D., Charbonnier, J., Parisse, C., Chakir, A., & Brion, J. (1995). Ozone UV spectroscopy. II. Absorption cross-sections and temperature dependence. *Journal of atmospheric chemistry*, 21, 263-273. <https://doi.org/10.1007/BF00696758>
- Park, J., Liu, X., Bak, J., & Chance, K. (2026). TEMPO Ozone Profile Retrieval Algorithm Theoretical Basis Document.
- Park, J., Liu, X., Houch, J., Haffner, D., & Chnace, K. (2026). Algorithm Description for the TEMPO Total Ozone Retrieval Algorithm.
- TEMPO Validation Team (2023). Tropospheric Emissions: Monitoring of Pollution (TEMPO) – Level 2 Science Data Product Validation Plan, SAO-DRD-11.
- Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Liu, X., Park, J., Petropavlovskikh, I., Hanisco, T.F., Hanisco, T., Szykman, J., Valin, L., Baumann, E., Cede, A., Tiefengraber, M., Gebetsberger, M., Uesato, I., Zheng, X., Ahn, S., Chang, L., Lee, W., Kim, J., Lee, H., Baek, K., Redondas, A., Fujiwara, M., Wang, T., Grutter, M., Houck, J., Haffner, D., & Lee, S. (2025). Geostationary Satellites Total Ozone Observations: First Results on Ground-based Networks Validation Efforts for TEMPO and GEMS. *Geophysical Research Letters*, 52, e2025GL114768. <https://doi.org/10.1029/2025GL114768>
- Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A., Hilton, B. B., Nicks, D. K., Newchurch, M. J., Carr, J. L. Janz, S. J., Andraschko, M. R., Arola, A., Baker, B. D., Canova, B. P., Chan Miller, C., Cohen, R. C., Davis, J. E., Dussault, M. E., Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter, M., Herman, J. R., Houck, J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C., Lindfors, A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R., O'Sullivan, E. J., Palmer, P. I., Pierce, R. B., Pippin, M. R., Saiz-Lopez, A., Spurr, R. J. D., Szykman, J. J., Torres, O., Veeffkind, J. P., Veihelmann, B., Wang, H., Wang, J., & Chance, K. (2017). Tropospheric emissions: Monitoring of pollution (TEMPO). *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 17-39. <https://doi.org/10.1016/j.jqsrt.2016.05.008>